

LETTERS TO THE EDITOR**ALTERNATIVE SAFER APPROACH THROUGH THE CAROTID ARTERY FOR TRANSCATHETER AORTIC VALVE IMPLANTATION****To the Editor:**

We read with deep interest the case report by Modine and coauthors¹ about the first transcatheter aortic valve implantation procedure through the left carotid artery access. The patient had a transient right hemiparesis after the procedure, probably because of the hypoperfusion of the left cerebral hemisphere despite the presence of a healthy circle of Willis or because of dissection of the left carotid artery.

We would like to mention a modification of the carotid access that addresses both complications. Direct visual introduction of a sheath through an open arteriotomy, then snaring the artery on the introducer and advancing the hydrophilic guidewire under fluoroscopic control would be helpful to prevent iatrogenic dissection. On the other hand, perfusion of the distal carotid artery with the aid of a centrifugal pump (eg, femoral artery to carotid artery) is an option to prevent the cerebral hypoperfusion. Our group successfully performed an experimental study for endovascular repair of the aortic arch with distal carotid perfusion.²

Open carotid arteriotomy access with distal perfusion would also be helpful to prevent the embolic shower, at least to the ipsilateral cerebral hemisphere, during the instrumentation,

balloon valvuloplasty, and implantation of the prosthesis. Allowing the free flow of the proximal carotid artery for a few seconds would clear off the debris.

We congratulate Modine and coauthors¹ for their novel application of the transcatheter aortic valve implantation procedure.

Harun Arbatli, MD^a

Oguz Yilmaz, MD^b

Mustafa Cikiricioglu, MD, PhD^c

^aDepartment of Cardiovascular Surgery

Maltepe University Faculty of Medicine

Istanbul, Turkey

^bDepartment of Cardiovascular Surgery

Memorial Hospital

Istanbul, Turkey

^cDivision of Cardiovascular Surgery

Department of Surgery

University Hospital of Geneva

Geneva, Switzerland

References

1. Modine T, Lemesle G, Azzaoui R, Sudre A. Aortic valve implantation with the CoreValve ReValving System via left carotid artery access: first case report. *J Thorac Cardiovasc Surg.* 2010;140:928-9.
2. Numan F, Arbatli H, Brzdzewski W, Cikiricioglu M. Total endovascular aortic arch reconstruction via fenestration in situ with cerebral circulatory support: an acute experimental study. *Interact Cardiovasc Thorac Surg.* 2008;7:535-8.

doi:10.1016/j.jtcvs.2010.10.052

Reply to the Editor:

We congratulate Arbatli and colleagues on their letter commenting on the carotid route used to achieve transcatheter aortic valve implantation. They give a reasonable explanation for the transient ischemic accident the patient experienced in our reported case. Indeed, hypoperfusion of the left cerebral hemisphere may have played a major role. This was probably mostly related to the carotid dissection that occurred initially. Arbatli and colleagues suggest ameliorating this technique by direct visualization of

sheath introduction through the open arteriotomy under fluoroscopic control. This precaution is now used in all our surgical transcatheter aortic valve implantations with carotid and subclavian arteries.¹ They also suggest, on the basis of an experimental study that they performed,² the use of distal perfusion centrifugal pumps. We do not recommend that technique. Indeed, the rationale behind transcatheter aortic valve implantation is to achieve minimally invasive implantation procedures. A complete study of circle of Willis perfusion, together with a full respect of the different steps to achieve this indication, could guarantee the success of this procedure. Indeed, without the occurrence of any complications, our group performed implantations in additional patients by this pathway. Finally Arbatli and colleagues point out the advantage of this surgical access as offering the possibility of preventing potential ipsilateral embolic shower during the balloon valvuloplasty and the valve implantation itself, which are the highest moments of cerebral embolic risks. Indeed, the carotid arteriotomy could be used as a window to clear off atherosclerotic debris during the procedure.

Transcatheter aortic valve implantation indications are growing. Exchanges of experiences are needed to optimize surgical implications and results.

Thomas Modine, MD

Georges Fayad, MD

Mohamad Koussa, MD

Service de Chirurgie

Cardiovasculaire

Hôpital Cardiologique

Centre Hospitalier Régional et

Universitaire de Lille

Lille, France

References

1. Modine T, Obadia JF, Choukroun E, Rioufoul G, Sudre A, Laborde JC, et al. Transcatheter aortic valve implantation using the axillary/subclavian access: Feasibility and early clinical outcomes. *J Thorac Cardiovasc Surg.* Epub 2010 Apr 28.

The Editor welcomes submissions for possible publication in the Letters to the Editor section that consist of commentary on an article published in the Journal or other relevant issues. Authors should: • Include no more than 500 words of text, three authors, and five references. • Type with double-spacing. • See <http://jtcvs.ctsnetjournals.org/misc/ifa.shtml> for detailed submission instructions. • Submit the letter electronically via jtcvs.editorialmanager.com. Letters commenting on an article published in the JTCVS will be considered if they are received within 6 weeks of the time the article was published. Authors of the article being commented on will be given an opportunity of offer a timely response (2 weeks) to the letter. Authors of letters will be notified that the letter has been received. Unpublished letters cannot be returned.